

MATHEMATICS

A NOTE ON THE ABSOLUTE BOUND FOR SYSTEMS OF LINES

BY

TOM H. KOORNWINDER

(Communicated by Prof. W. T. VAN EST at the meeting of September 27, 1975)

ABSTRACT

The absolute bound for the cardinality of sets of lines in \mathbb{R}^n or \mathbb{C}^n having a prescribed number of angles, which was first obtained by Delsarte, Goethals & Seidel, is derived in a new, very quick way.

One of the main results in DELSARTE, GOETHALS & SEIDEL [1] is the derivation of an absolute bound for the cardinality of sets of lines having a prescribed number of angles, both in real and in complex Euclidean n -space. In order to formulate this result (cf. Theorem 1) we first introduce some notation.

Let (\cdot, \cdot) denote the real or Hermitian inner product in \mathbb{R}^n or \mathbb{C}^n , respectively. Let Ω_n be the unit sphere in \mathbb{R}^n or \mathbb{C}^n . Let x_1, x_2, \dots, x_n be real (or complex) coordinates on \mathbb{R}^n (or \mathbb{C}^n). In the case \mathbb{R}^n , let $\text{Hom}(l)$ be the class of functions on Ω_n which are restrictions to Ω_n of homogeneous polynomials of degree l in x_1, x_2, \dots, x_n . In the case \mathbb{C}^n , let $\text{Hom}(l, k)$ be the class of functions on Ω_n which are restrictions to Ω_n of polynomials $F(x_1, \dots, x_n; \bar{x}_1, \dots, \bar{x}_n)$, homogeneous of degree l in x_1, \dots, x_n and homogeneous of degree k in $\bar{x}_1, \dots, \bar{x}_n$ (cf. KOORNWINDER [2, § 2]). Then

$$(1) \quad M(l) = \dim \text{Hom}(l) = \binom{n+l-1}{n-1},$$

$$(2) \quad M(l, k) = \dim \text{Hom}(l, k) = \binom{n+l-1}{n-1} \binom{n+k-1}{n-1}.$$

Let $A = \{\alpha_1, \alpha_2, \dots, \alpha_s\}$ be a subset of the interval $[0, 1)$. Put $\varepsilon = 1$ if $0 \in A$ and $\varepsilon = 0$ if $0 \notin A$. Let X be a finite subset of Ω_n of cardinality $|X|$ such that $|(\xi, \eta)|^2 \in A$ if $\xi, \eta \in X$ and $\xi \neq \eta$.

THEOREM 1. $|X| \leq M(2s - \varepsilon)$ in the case \mathbb{R}^n , and $|X| \leq M(s, s - \varepsilon)$ in the case \mathbb{C}^n .

This theorem was proved in [1, Theorem 6.1] by using the decomposition of functions on Ω_n in terms of real or complex spherical harmonics and

by expressing the zonal spherical harmonics on Ω_n as Jacobi polynomials. Below we give a new, almost trivial proof of Theorem 1.

PROOF OF THEOREM 1. For each $\xi \in X$ define the polynomial

$$(3) \quad F_{\xi}(x) = ((x, \xi))^s \prod_{\alpha \in A \setminus \{0\}} \frac{|(x, \xi)|^2 - \alpha(x, x)}{1 - \alpha}.$$

Then the restriction of F_{ξ} to Ω_n belongs to $\text{Hom}(2s - \varepsilon)$ [in the case \mathbb{R}^n and to $\text{Hom}(s, s - \varepsilon)$ in the case \mathbb{C}^n]. Furthermore

$$F_{\xi}(\eta) = \delta(\xi, \eta), \quad \xi, \eta \in X.$$

Hence the functions F_{ξ} , $\xi \in X$, are linearly independent. It follows that $|X|$ is bounded by the dimension of $\text{Hom}(2s - \varepsilon)$ or $\text{Hom}(s, s - \varepsilon)$. \square

*Mathematisch Centrum
2e Boerhaavestraat 49
Amsterdam-1005, The Netherlands*

REFERENCES

1. DELSARTE, P., J. M. GOETHALS & J. J. SEIDEL, Bounds for systems of lines and Jacobi polynomials, Philips Res. Repts. 30, 91*-105* (1975).
2. KOORNWINDER, T. H., The addition formula for Jacobi polynomials, II, The Laplace type integral representation and the product formula, Math. Centrum, Amsterdam, Report TW 133 (1972).